

A CANCELLATION THEOREM FOR IDEALS

CRAIG HUNEKE

ABSTRACT. We prove cancellation theorems for special ideals in Gorenstein local rings. These theorems take the form that if $KI \subseteq JI$, then $K \subseteq J$.

1. INTRODUCTION

One of the basic themes in commutative Noetherian ring theory is that of cancellation of ideals: if I, J, K are ideals in a commutative ring R and $IK = JK$, then one cannot in general conclude that $I = J$. Of course for Dedekind domains it is an old and famous theorem that one can do so. In general, one must be satisfied with ‘cancellation up to integral closures’, i.e. in concluding that the integral closure of I is equal to the integral closure of J .

It is of considerable interest to understand when one can cancel: rarely is what experience teaches. However, the point of this paper is to prove a cancellation theorem for special ideals. This cancellation has a number of immediate corollaries, e.g. to theorems of Briançon-Skoda type which say that the integral closure of a certain power of I is contained in I , as well as a number of corollaries dealing with reduction numbers and the syzygetic property.

We recall relevant definitions:

Definition 1.1. Let R be a ring and let I be an ideal of R . An element $x \in R$ is *integral over I* if x satisfies an equation of the form $x^n + a_1x^{n-1} + \cdots + a_n = 0$, where $a_j \in I^j$ for $1 \leq j \leq n$. The *integral closure* of I , denoted by $\bar{I}$, is the set of all elements integral over I . This set is an ideal.

Closely related to integral closure is the idea of a reduction. Recall that $J \subseteq I$ is said to be a *reduction* of I if there exists an integer $n \geq 0$ such that $I^{n+1} = JI^n$. The least such n is said to be the reduction number of I with respect to J . It is denoted $r_J(I)$. If J is a reduction of I and properly contains no other reduction of I , J is said to be a *minimal reduction*.

The *analytic spread* of an ideal I in a local Noetherian ring $(R, \mathfrak{m})$ with infinite residue field is the least number of generators of any minimal reduction of I . Equivalently, it is

1991 *Mathematics Subject Classification.* Primary 13D40, 13A30, 13H10.

The author is grateful to the Max-Planck Institutue for its support while visiting during Fall 1998, where part of this work was completed. The author also thanks the NSF for partial support.

the Krull dimension of the ‘fiber cone’, $R[It]/\mathbf{m}R[It]$. The *analytic deviation* of I , denoted $\text{ad}(I)$, is the analytic spread of I minus the height of I .

Our main theorem is Theorem 2.2 below. Its most basic case says that if R is a regular local ring, P a prime of R with $\dim(R/P) = 1$, and J any minimal reduction of P , then $IP \subseteq JP$ implies $I \subseteq J$ for an arbitrary ideal I .

2. MAIN RESULT

We begin with an elementary lemma.

Lemma 2.1. *Let $(R, \mathbf{m})$ be a local Gorenstein ring. Let $I \subseteq R$ be an ideal of height g . Set $d = \dim(R/I)$. Let $\underline{a} = (a_1, \dots, a_g)$ be a regular sequence in I . Then $R/(\underline{a} : I)$ satisfies Serre’s condition S_2 iff $H_{\mathbf{m}}^{d-1}(R/I) = 0$.*

Proof. We reduce first to the case in which I is unmixed. Let K be the unmixed part of the primary decomposition of I , so that K/I has dimension strictly less than d . The long exact sequence on local cohomology applied to the short exact sequence,

$$0 \longrightarrow K/I \longrightarrow R/I \longrightarrow R/K \longrightarrow 0$$

shows that $H_{\mathbf{m}}^{d-1}(R/K) = 0$ as it lies between $H_{\mathbf{m}}^{d-1}(R/I)$, which is 0 by assumption, and $H_{\mathbf{m}}^d(K/I)$, which is 0 since d is greater than the dimension of K/I . We can replace I by K if we prove that $(\underline{a} : I) = (\underline{a} : K)$. It suffices to check this at the associated primes of $(\underline{a} : K)$ (as $I \subseteq K$ implies that $(\underline{a} : K) \subseteq (\underline{a} : I)$), and all of these are associated to $\underline{a}$. If Q is such a prime and $I \not\subseteq Q$, then $(\underline{a})_Q = (\underline{a} : I)_Q = (\underline{a} : K)_Q$. If $I \subseteq Q$, then $I_Q = K_Q$, and again $(\underline{a} : I)_Q = (\underline{a} : K)_Q$. We replace I by K and assume it is unmixed.

Assume that $H_{\mathbf{m}}^{d-1}(R/I) = 0$. Let Q be a prime containing $\underline{a} : I$, and assume that the height of Q is at least $g + 1$. If Q does not contain I , then $(\underline{a} : I)_Q = (\underline{a})_Q$ and the result follows. Henceforth we assume $I \subseteq Q$.

Choose an element $t \in Q$ which is not a zero-divisor on either R/I or $R/(\underline{a}_1, \dots, \underline{a}_g)$. Apply $\text{Hom}_R(R/I, _)$ to the short exact sequence $0 \longrightarrow R \xrightarrow{t} R \longrightarrow S \longrightarrow 0$, where $S = R/Rt$. Note that $\text{Ext}_R^{g+1}(R/I, R)$ is the Matlis dual of $H_{\mathbf{m}}^{d-1}(R/I)$ (recalling R is Gorenstein), and hence is zero. Thus we get a short exact sequence,

$$0 \longrightarrow \text{Ext}_R^g(R/I, R) \xrightarrow{t} \text{Ext}_R^g(R/I, R) \longrightarrow \text{Ext}_R^g(R/I, S) \longrightarrow 0.$$

Since $a_1, \dots, a_g$ are a regular sequence on R and R/tR in I , we may identify this last exact sequence with the sequence,

$$0 \longrightarrow \text{Hom}_R(R/I, R/\underline{a}) \xrightarrow{t} \text{Hom}_R(R/I, R/\underline{a}) \longrightarrow \text{Hom}_R(R/I, S/\underline{a}S) \longrightarrow 0.$$

This gives the identity,

$$(1) \quad (\underline{a} : I) + tR = (\underline{a} + Rt : I).$$

In particular, since $a_1, \dots, a_g, t$ form a regular sequence, the ideal $(\underline{a} : I) + tR$ is unmixed, necessarily of height $g + 1$. But then $\text{depth}((R/((a_1, \dots, a_g) : I))_Q) \geq 2$, unless Q has height exactly $g + 1$, in which case this localization is Cohen-Macaulay. Thus $R/(\underline{a} : I)$ satisfies S_2 .

Conversely, suppose that $R/(\underline{a} : I)$ satisfies S_2 . Notice that $\text{Ext}_R^{g+1}(R/I, R)$ has an annihilator of height at least $g + 2$ since at all primes of height $g + 1$, after localization this module is zero as R/I is S_1 . Choose an element t , a zero-divisor on both R/I or $R/(a_1, \dots, a_g)$ such that $t\text{Ext}_R^{g+1}(R/I, R) = 0$. The same exact sequences as above then give the isomorphism

$$((\underline{a} + Rt) : I)/((\underline{a} : I) + tR) \cong \text{Ext}_R^{g+1}(R/I, R).$$

This proves that the associated primes of $\text{Ext}_R^{g+1}(R/I, R)$ are contained in the associated primes of $R/((\underline{a} : I) + tR)$, and since $R/(\underline{a} : I)$ satisfies S_2 , all such primes have height at most $g + 1$. It follows that $\text{Ext}_R^{g+1}(R/I, R) = 0$. $\square$

Theorem 2.2 (Cancellation). *Let $(R, \mathfrak{m})$ be a local Gorenstein ring. Let $I \subseteq R$ be an unmixed ideal of height g , and let $a_1, \dots, a_{g+1} \in I$. Assume that $a_1, \dots, a_g$ form a regular sequence which generates I locally at the localization of every minimal of I , and that a_{g+1} is not contained in any minimal prime of $(a_1, \dots, a_g)$ which does not contain I . Set $J = (a_1, \dots, a_g, a_{g+1})$. Finally assume that $H_{\mathfrak{m}}^{d-1}(R/I) = 0$ (e.g. if R/I is Cohen-Macaulay), where $d = \dim(R/I)$. If $KI \subseteq JI$, then $K \subseteq J$.*

Proof. Observe that the assumptions of the theorem force $\underline{a} : I = \underline{a} : a_{g+1}$. We first reduce to the situation in which R/I has dimension one; this is the main case. Our assumption on the local cohomology is made to reduce to this case.

To prove $K \subseteq J$, it suffices to prove $K_Q \subseteq J_Q$ for every associated prime Q of J . We claim all such associated primes have height at most $g + 1$. This will then reduce to the case in which R/I has dimension one by localization. Let Q be an associated prime of J and suppose that Q has height at least $g + 2$. The exact sequence

$$0 \longrightarrow R/((a_1, \dots, a_g) : a_{g+1}) \longrightarrow R/(a_1, \dots, a_g) \longrightarrow R/J \longrightarrow 0$$

induced by multiplication by a_{g+1} on the first map shows that Q must be associated either to $R/(a_1, \dots, a_g)$, which is impossible, or $\text{depth}((R/((a_1, \dots, a_g) : a_{g+1}))_Q) \leq 1$. By Lemma 2.1, $R/((a_1, \dots, a_g) : a_{g+1})$ is S_2 . As Q has height at least $g + 2$, this is a contradiction, proving our claim.

We have reduced to the case in which $\dim(R/I) = 1$. Choose an element s , not a zero-divisor on R/I , such that $\underline{a} : s = I$. This is possible since I is generically equal to $\underline{a}$ and is unmixed: specifically choose an element s which is in all the primary components of the ideal $\underline{a}$ whose radicals do not contain I and in addition choose s not in any associated prime of I . As both I and $\underline{a}$ are unmixed this is possible using prime avoidance. The condition that s not be a zero-divisor on R/I is immediate from the choice made. We can write $\underline{a} = N \cap I$, where the minimal primes over N do not contain I . We have chosen $s \in N$ but not a zero-divisor modulo I . Hence $\underline{a} : s = (N : s) \cap (I : s) = I$. Set $b = a_{g+1} + s$, and

set $\mathfrak{A} = (a_1, \dots, a_g, b)$. Observe that $\mathfrak{A}$ is $\mathbf{m}$ -primary, as b is not contained in the union of the minimal primes over $\underline{a}$.

Set $q = (\underline{a} : I) + I$. We claim that $K \subseteq \mathfrak{A} : q$. Clearly $K(\underline{a} : I) \subseteq \underline{a} \subseteq \mathfrak{A}$, since $K \subseteq I$. This last containment follows since it is enough to check it at the localizations of the associated primes of I , which are all minimal over I . After localizing at such a prime, I is equal to $\underline{a}$, and our assumption then reads (after localization at a minimal prime Q) that $K_Q(\underline{a})_Q \subseteq (\underline{a})_Q^2$. As $\underline{a}$ is generated by a regular sequence it then follows that $K_Q \subseteq (\underline{a})_Q$. Hence $K \subseteq I$. To prove that $KI \subseteq \mathfrak{A}$, it is enough to prove that $JI \subseteq \mathfrak{A}$ by our basic assumption. As $\underline{a}I \subseteq \mathfrak{A}$, we only need to prove that $a_{g+1}I \subseteq \mathfrak{A}$. But for $i \in I$, we may write $a_{g+1}i = (a_{g+1} + s)i - si \in \mathfrak{A}$ since $si \in \underline{a}$.

We next claim that $\mathfrak{A} : s = q$. One direction is much the same as above. We know $sI \subseteq \underline{a}$ by choice of s . If $t \in \underline{a} : I$, then we can write $ts = tb - ta_{g+1} \in \mathfrak{A}$ since $ta_{g+1} \in \underline{a}$. Thus $q \subseteq \mathfrak{A} : s$. Conversely, suppose that $su \in \mathfrak{A}$. Write $su = bv + w$, where $w \in \underline{a}$. Then $s(u - v) \in J \subseteq I$, and thus $u - v \in I$. Note that $Ibv \subseteq Isu + Iw \subseteq \underline{a}$. Since b is a regular element modulo $\underline{a}$, we see that $Iv \subseteq \underline{a}$. Thus $u = (u - v) + v \in I + \underline{a} : I = q$.

As $\mathfrak{A}$ is an $\mathbf{m}$ -primary ideal in a Gorenstein local ring, generated by a regular sequence, we know that $K \subseteq \mathfrak{A} : q = \mathfrak{A} : \mathfrak{A} : s = \mathfrak{A} + Rs = J + Rs$. Hence $K \subseteq (J + Rs) \cap I = J + (I \cap Rs) \subseteq J + Is \subseteq J$, finishing the proof. $\square$

Corollary 2.3. *Let $(R, \mathbf{m})$ be a local Gorenstein ring with infinite residue field and suppose that I is an unmixed ideal, generically a complete intersection, such that the $ad(I) = 1$. Assume that $H_{\mathbf{m}}^{d-1}(R/I) = 0$ where d is the Krull dimension of R/I . Let J be an arbitrary minimal reduction of I . Then $r_J(I) \neq 1$. In particular, if R is a regular ring and P is a prime with $\dim(R/P) = 1$, then P cannot have reduction number 1 with respect to a minimal reduction of P .*

Proof. The last statement follows immediately from the first. The assumptions guarantee that a given minimal reduction J of I is generated by $g + 1$ elements which can be chosen as in the conditions of Theorem 2.2. If the reduction number is one, then $P^2 = JP$, and cancellation then proves that $P \subseteq J$, that is $P = J$, and the reduction number is 0. $\square$

Theorem 2.4. *Let $(R, \mathbf{m})$ be a regular local ring with infinite residue field. Suppose that I is an unmixed ideal, generically a complete intersection, such that the $ad(I) = 1$. Assume that $H_{\mathbf{m}}^{d-1}(R/I) = 0$ where d is the Krull dimension of R/I . Set $g = \text{height}(I)$. Then $\overline{I^g} \subseteq J$, where J is an arbitrary minimal reduction of I . In particular, if P is a prime of R with $\dim(R/P) = 1$, then $\overline{P^g} \subseteq J$, for an arbitrary minimal reduction J of P .*

Proof. By the results in [AH] and [L], we know that

$$\overline{I^{g+1}} \subseteq JI.$$

Since $I(\overline{I^g}) \subseteq \overline{I^{g+1}}$, the result follows at once from the cancellation theorem. $\square$

The above result suggests at least the possibility that for every prime P in a regular local ring R , $\overline{P^g} \subseteq J$, for $g = \text{height}(P)$ and J an arbitrary minimal reduction of P . This would be a considerable improvement on the usual Briançon-Skoda results which

only give $\overline{P^d} \subseteq J$ in general, for $d = \dim(R)$. Trying to find an example in which the stronger claim $\overline{P^g} \subseteq J$ does not hold turned out to be difficult, and no counterexample was found. In fact, nearly every example checked, no matter what the analytic spread of P , had the containment $P^2 \subseteq J$. Example 2.7 below gives a height three prime ideal such that $P^2 \not\subseteq J$. The examples below deal with powers, rather than the integral closure of powers as this author does not know a good way to compute the integral closures of the powers of these ideals. The examples were all computed on MACAULAY, in the highest characteristic allowable.

Example 2.5. Let P be the defining ideal of $k[t^4 + s^4, t^3s^2, t^2s^4 + ts^5, ts^6]$ in the ring $R = k[X_1, \dots, X_4]$. Then P is height two, is generated by 5 elements, say $a_1, \dots, a_5$, and R/P is not Cohen-Macaulay. The fiber cone $S/\mathbf{m}S$, where $\mathbf{m} = (X_1, \dots, X_4)$ and $S = R[Pt]$, is isomorphic to $k[T_1, \dots, T_5]/(T_1T_5)$. It follows that the analytic spread of P is 4. A minimal reduction is $J = (a_1 + a_5, a_2, a_3, a_4)$. One can check that $P^2 \subseteq J$.

Example 2.6. For a slightly more complicated example, consider the defining ideal of the ring $k[s^3, t^3, u^3, s^2t + stu, st^2 + tu^2]$. This ideal P is height two, generated by 8 elements. If we map $R[A, B, C, D, E, F, G, H]$ onto the Rees algebra using these 8 generators, the fiber cone $k[A, B, C, D, E, F, G, H]/I$ is defined by $I = (F^2 - 1/2DG + EG - FG, EF - 1/2DG - 2EG - FG + G^2, E^2 + 4/9CG + 5/6DG + EG + FG - G^2 + 1/18EH - 1/18FH - 1/6GHDF - CG, DE - 1/9CG - 1/3DG + 1/9EH - 1/9FH - 1/3GH, D^2 + 16/9CG - 2/3DG + 2/9EH - 2/9FH + 4/3GH, CF + GH, CE - 1/3CG + 1/3EH - 1/3FH, CD - 2/3CG + 2/3EH + 4/3FH - 2GH, C^2 + DH)$. This ideal has height four, so the analytic spread of P is $4 = 8 - 4$. A minimal reduction J is given by the first three generators corresponding to A, B, C , together with the sum of all the rest of the specified generators. Precisely, if we map $k[a, b, c, d, e]$ onto $k[s^3, t^3, u^3, s^2t + stu, st^2 + tu^2]$, then J is generated by the four elements,

$$f_1 = a^2bc - bc^3 + 3abcd - 3bcd^2 + 3abd^2 - 3bcd^2 - bd^3 - cd^3 - 3a^2be - 3abce - 3abde - 3d^3e + 3ade^2 + 3d^2e^2 + ae^3 + ce^3,$$

$$f_2 = a^2b^2 + 1/2abc^2 - 1/2bc^3 - 3/2bc^2d - 3/2bcd^2 - 1/2bd^3 - 3abce - 3/2abde + 3/2d^2e^2 - 1/2ae^3 + 1/2ce^3,$$

$$f_3 = abd^3 + c^2d^3 - a^2bde - abcde - 3abd^2e - 2d^4e - a^2be^2 - abce^2 - 3acde^2 + 3ad^2e^2 + d^3e^2 - a^2e^3 - ace^3 + 3ade^3, \text{ and}$$

$$f_4 = ab^2c^2 + b^2c^3 + ab^2cd + abc^2d + 3b^2c^2d + 2bc^3d + a^2bd^2 + 2ab^2d^2 + abcd^2 + 3b^2cd^2 + 5bc^2d^2 + b^2d^3 + 3bcd^3 - 3c^2d^3 + bd^4 - 3cd^4 - d^5 - a^3be - 3ab^2ce + 4abc^2e - 4a^2bde - 7ab^2de + 2abcde + 2bc^2de + ad^3e - 3bd^3e - cd^3e + d^4e + 3a^2be^2 + 2ab^2e^2 - 6abce^2 + 2bc^2e^2 + 3abde^2 + 9acde^2 + 6bcde^2 + 3bd^2e^2 + 3a^2e^3 + 5abe^3 + 3ace^3 - bce^3 - 4ade^3 - 2cde^3 + d^2e^3 - 3ae^4 - 2de^4 - 2e^5.$$

One can check that $P^2 \subseteq J$.

Example 2.7. Let I be the ideal of maximal minors of the 4 by 6 matrix,

$$A = \begin{pmatrix} a & b & c & e & f & d \\ d & a & b & c & e & f \\ e & d & a & b & c & e \\ f & e & d & a & b & c \end{pmatrix}.$$

The ideal I has generic height in $R = k[a, b, c, d, e, f]$, namely $3 = 6 - 4 + 1$. In particular R/I is Cohen-Macaulay. This ideal was chosen so that after the specialization $d = e = f = 0$, the specialized ideal becomes the 4th power of (a, b, c) , and the square of this ideal is not contained in any minimal reduction of it. One can check (with a huge amount of computer time!) that I is reduced, and has analytic spread 6. We computed a particular minimal reduction J (which is extremely complicated), and for this particular reduction, $P^2 \not\subseteq J$. However, $P^2 \subseteq J : \mathbf{m}$, where $\mathbf{m}$ is the obvious maximal ideal. In particular $P^3 \subseteq J$. The calculation of this example took more than a week of computer time, and discouraged the author from looking much further!

We recall some concepts before stating the next theorem. Let R be a Noetherian ring and I an ideal. If I is generated by $a_1, \dots, a_n$ we can map $S = R[T_1, \dots, T_n]$ onto $R[It]$ by sending T_i to $a_i t$. The kernel of this map we denote by Q . This ideal is graded, and by Q_j we denote the j th graded piece of Q . The ideal I is said to be *syzygetic* if $Q_2 \subseteq S_1 Q_1$. See [V, 2.1] for a discussion of this concept. Some known classes of syzygetic ideals are of course those of linear type, where $SQ_1 = Q$, but also include, e.g., all height two reduced ideals in a 3-dimensional regular local ring.

Theorem 2.8. *Let R be a Gorenstein local ring and P a prime ideal such that $ad(P) = 1$, R_P is regular, and $H_{\mathbf{m}}^{d-1}(R/P) = 0$ where d is the Krull dimension of R/P . If $\mu(P) = g+2$, where $g = \text{height}(P)$, then P is syzygetic.*

Proof. Suppose not. Fix a generating set $a_1, \dots, a_{g+2}$ of P . Then there is a homogeneous polynomial of degree two, $H \in R[T_1, \dots, T_{g+2}]$ such that $H(a_1, \dots, a_{g+2}) = 0$ and such that $H \notin \mathfrak{A}$, where $\mathfrak{A} = SQ_1$ is the ideal defining the symmetric algebra of P , i.e. the ideal in $R[T_1, \dots, T_{g+2}]$ generated by all linear polynomials which vanish at $a_1, \dots, a_{g+2}$. After possibly changing the generators, we may assume $H = rT_1^2 + P$, where P is a polynomial in the ideal $(T_2, \dots, T_{g+2})$; we may further assume that $J = (a_2, \dots, a_{g+2})$ is a minimal reduction of P , and that P is generically generated by $a_2, \dots, a_{g+1}$ which form a regular sequence. But then $ra_1^2 \in JI$, and hence $rI^2 \subseteq JI$. By the cancellation theorem, $rI \subseteq J$, so there is a linear polynomial $L = rT_1 + L'$, where $L' \in (T_2, \dots, T_{g+2})$, such that $L(a_1, \dots, a_{g+2}) = 0$. Consider the polynomial $H - T_1 L = G$. The polynomial G can be written in the form

$$G = T_2 L_2 + \dots + T_{g+2} L_{g+2}$$

for linear polynomials L_i . Set $l_i = L_i(a_1, \dots, a_{g+2})$. Then $l_{g+2} \in (a_2, \dots, a_{g+1}) : a_{g+2} \cap P$. Since $a_2, \dots, a_{g+1}$ generically generate P this intersection is just $(a_2, \dots, a_{g+1})$. Hence there is another linear polynomial $L^* = r_2 T_2 + \dots + r_{g+1} T_{g+1}$ such that $l_{g+2} = L^*(a_2, \dots, a_{g+1})$. In other words $L_{g+2} - L^*$ is a linear polynomial in $\mathfrak{A}$. Modulo $\mathfrak{A}$ we can write $G = T_2 F_2 + \dots + T_{g+1} F_{g+1}$ for some new choice of linear polynomials F_i . But then $G \in Q_2 \cap (T_2, \dots, T_{g+1})$, and $a_2, \dots, a_{g+1}$ form a regular sequence. The next lemma then finishes the proof. It is well-known to experts, but a suitable reference was not found. (See however [AH, (2.5)].)

Lemma 2.9. *Let $(R, \mathbf{m})$ be a Noetherian local ring and let I be an ideal. Assume that $(a_1, \dots, a_n, a_{n+1}, \dots, a_m) = I$, and that $\underline{a} = a_1, \dots, a_n$ form a regular sequence. Let $R[T_1, \dots, T_m] \longrightarrow R[It]$ be the surjection of the polynomial ring $R[T_1, \dots, T_m]$ onto $R[It]$*

sending T_i to $a_i t$. Let Q be the kernel of this surjection with the obvious grading. Then $(T_1, \dots, T_n) \cap Q_2 \subseteq S_1 Q_1$.

Proof. By descending induction on i , we prove that

$$(T_1, \dots, T_i) \cap Q_2 \subseteq S_1 Q_1 + (T_1, \dots, T_{i-1}) \cap Q_2.$$

Since $(T_1) \cap Q_2 = T_1 Q_1$ as a_1 is a nonzerodivisor, the Lemma will follow.

Suppose that $G \in (T_1, \dots, T_i) \cap Q_2$, and write $G = \sum_{j=1}^i T_j G_j$, with G_j linear forms. Write $g_j = G_j(a_1, \dots, a_m)$. Then $g_i \in (a_1, \dots, a_{i-1}) : a_i = (a_1, \dots, a_{i-1})$, so we may write $g_i = \sum_{k=1}^{i-1} a_k r_k$. Set $H = \sum_{k=1}^{i-1} T_k r_k$, so that $G_i - H \in Q_1$. Then $G = \sum_{j=1}^{i-1} T_j G_j + T_i(G_i - H) + T_i H \in (T_1, \dots, T_{i-1}) + S_1 Q_1$, from which the result follows. $\square$

Theorem 2.10. *Let I be an unmixed ideal in a regular local ring of dimension d . such that I_P is generated by a regular sequence of length $g < d$ for every prime $I \subseteq P$ with $P \neq \mathfrak{m}$. Then $\overline{I^{d-1}} \subseteq J$, for every reduction J of I .*

Proof. We induce upon the dimension of R/I . If the dimension is 1, then Theorem 2.4 gives us the conclusion. Choose a regular sequence $\underline{a} = a_1, \dots, a_g \in I$ which generated I_P for every minimal prime P above I . Set $K = \underline{a} : I + I$. Then the dimension of R/K is strictly less than the dimension of R/I . We claim that K is unmixed and K_Q is generated by a regular sequence of length $g + 1$ for every prime $K \subseteq Q$ with $Q \neq \mathfrak{m}$. This follows from the next Lemma:

Lemma 2.11. *Let I be an unmixed ideal in a local Cohen-Macaulay ring $(R, \mathfrak{m})$, and assume that I is generically generated by a regular sequence of length g . Let $\underline{a} = a_1, \dots, a_g$ be a regular sequence in I which generates I locally at all minimal primes of I . Define an ideal $K = (\underline{a} : I) + I$. Then K is an unmixed ideal of height $g + 1$. If I_P is generated by a regular sequence of length g for all primes $I \subseteq P$ such that $\dim((R/I)_P) \leq 1$, then K is generically a complete intersection. More generally if $K \subseteq Q$ and I_Q is generated by a regular sequence, then K_Q is also generated by a regular sequence.*

Proof. We first prove that the height of K is at least $g + 1$. If P contains K and has height at most g , then P must contain I (hence P has height exactly g) and $(\underline{a})_P = I_P$ by assumption. Then $\underline{a} : I \not\subseteq P$, a contradiction. Hence $\text{height}(K) \geq g + 1$. Let Q be an associated prime of K . The exact sequence

$$0 \longrightarrow R/((\underline{a} : I) \cap I) \longrightarrow R/I \oplus R/(\underline{a} : I) \longrightarrow R/K \longrightarrow 0$$

shows that either Q is associated to I , to $\underline{a} : I$, or else $\text{depth}(R/((\underline{a} : I) \cap I)) \leq 1$. Both I and $\underline{a}$ are unmixed, so every associated prime of either I or $\underline{a} : I$ has height g . On the other hand, since $K((\underline{a} : I) \cap I) \subseteq \underline{a}$, and $\text{height}(K) \geq 1$, it follows that $(\underline{a} : I) \cap I = \underline{a}$. Then the height of Q is at most $g + 1$, which proves that K is unmixed of height $g + 1$.

The penultimate statement follows at once from the more general last statement of the theorem. Let Q be a prime containing K , such that I_Q is a complete intersection, say generated by $b_1, \dots, b_g$. Then $\underline{a} : (b_1, \dots, b_g)$ is generated by $a_1, \dots, a_g, \delta$, where δ is the

determinant of a transition matrix D writing the a_i in terms of the b_i . See [BH,(2.3.10)]. It follows that $K_Q = (a_1, \dots, a_g, \delta)_Q + (b_1, \dots, b_g)_Q = (b_1, \dots, b_g, \delta)_Q$ is generated by $g + 1$ elements, which are necessarily a regular sequence as the height of K is $g + 1$ and R is Cohen-Macaulay.

We can apply the induction to conclude that $\overline{K^{d-1}} \subseteq L$, for every reduction L of K . Let J be an arbitrary reduction of I . Then $J + \underline{a} : I$ is certainly a reduction of K , so we obtain that

$$\overline{I^{d-1}} \subseteq \overline{K^{d-1}} \cap I \subseteq (J + \underline{a} : I) \cap I = J + (\underline{a} : I \cap I) = J$$

which finishes the proof. $\square$

Another case is easy:

Theorem 2.12. *Let I be an ideal in a regular local ring $(R, \mathfrak{m})$ of dimension d . Assume that I is generically a complete intersection, and that R/I is Cohen-Macaulay of positive dimension. Then $\overline{I^{d-1}} \subseteq J$ for every reduction J of I .*

Proof. Induce upon the dimension of R/I . If this dimension is 1, we obtain the conclusion from Theorem 2.4. Assume it is bigger than 1. Denote the height of I by g . The set of primes q of height $g + 1$ such that I_q is not a complete intersection is finite in number, say $q_1, \dots, q_n$. Choose an element $y \in \mathfrak{m}$ a nonzerodivisor on R/I , and such that y is not contained in the union of $q_1, \dots, q_n$. Such a choice is possible by prime avoidance. Set $K = I + Ry$. Then K is generically a complete intersection, since if Q is minimal over K then $Q \neq q_i$ for any i , and hence I_Q is generated by g elements and K_Q by $g + 1$ elements. Clearly R/K is Cohen-Macaulay. Fix a reduction J of I . Then $J + Ry$ is a reduction of K . The induction implies that

$$\overline{I^{d-1}} \subseteq \overline{K^{d-1}} \cap I \subseteq (J + Ry).$$

We may replace y by y^n and use the Krull Intersection Theorem to prove that $\overline{I^{d-1}} \subseteq J$. $\square$

We end by giving one other somewhat surprising corollary of our main theorem.

Corollary 2.13. *Let $(R, \mathfrak{m})$ be a local Gorenstein ring. Let $I \subseteq R$ be an unmixed ideal of height g , and let $a_1, \dots, a_{g+1} \in I$. Assume that $a_1, \dots, a_g$ form a regular sequence which generates I locally at the localization of every minimal of I , and that a_{g+1} is not contained in any minimal prime of $(a_1, \dots, a_g)$ which does not contain I . Set $J = (a_1, \dots, a_g, a_{g+1})$. Finally assume that $H_{\mathfrak{m}}^{d-1}(R/I) = 0$ (e.g. if R/I is Cohen-Macaulay), where $d = \dim(R/I)$ and that the radical of J is equal to the radical of I (e.g. if J is a reduction of I). Then $I^n \subseteq J$ iff $I^{n+1} \cap J \subseteq JI$.*

Proof. One direction is obvious. If $I^n \subseteq J$, then $I^{n+1} \subseteq I^n \cdot I \subseteq JI$. For the other direction we assume that $I^{n+1} \cap J \subseteq JI$ and prove by descending induction on N that $I^N \subseteq J$. When $N = n$ we are done. By assumption, some power of I is contained in J . Suppose that $I^N \subseteq J$ and $N > n$. Then $I^N \subseteq I^{n+1} \cap J \subseteq JI$, and by Theorem 2.2, we may cancel the I to obtain that $I^{N-1} \subseteq J$. $\square$

Acknowledgement. The author thanks Ian Aberbach for many valuable conversations concerning the material in this paper.

Bibliography

- [AH] I. M. Aberbach and C. Huneke, *An improved Briançon-Skoda theorem with applications to the Cohen-Macaulayness of Rees rings*, Math. Ann. **297** (1993), 343–369.
- [BH] W. Bruns and J. Herzog, *Cohen-Macaulay Rings*, vol. 39, Cambridge studies in advanced mathematics, 1993.
- [L] J. Lipman, *Adjoints of ideals in regular local rings*, Math. Res. Letters **1** (1994), 1–17.
- [V] W. V. Vasconcelos, *Arithmetic of Blowup Algebras*, Cambridge University Press, Cambridge, 1994.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KANSAS, LAWRENCE, KS 66045
E-mail address: `huneke@math.ukans.edu`